



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Photic sense organs.—GUTTENBERG¹⁴ has demonstrated that two of the ombrophilous species of his local flora have a photo-sensitive epithelium, whose response consists in maintaining the leaf in the transverse heliotropic position. The mechanism is essentially the same as was found by HABERLANDT in the so-called velvet leaves, so abundant among the ombrophilous species of the tropical hydrophytic forests. The epidermal cells function as converging lenses, so that the protoplasmic membrane which covers the floor of the cell is not uniformly illuminated. In HABERLANDT's studies the bright spot was centrally located, but GUTTENBERG finds that for his species that it is excentric, because the papillosity is not centrally located. The result is the same in both cases, for the leaf is attuned to the distribution of interior illumination which exists when the leaf is in the transverse position. Actual tests showed that the petiole is not a factor in securing this position. Curiously enough the leaf assumes the horizontal position in diffuse light, such as occurs under the open sky on a cloudy day. In this case, however, the internal distribution of light is the reverse of that which exists under parallel rays, the central area of the floor wall being dark with a margin of increasing brightness. The stimulus apparently consists in an unequal illumination of the cell lumen.—RAYMOND H. POND.

Nature of chromatophores.—MERESCHKOWSKY¹⁵ holds that these bodies are not organs of the plant cell and never have been, but are foreign organisms which penetrated into the colorless plasma of the cells and live there as symbionts. In support of this notion he adduces the facts that the chromatophores multiply continuously by division and do not arise *de novo*; that they are in high degree independent of the nucleus; that they are completely analogous with zoochlorellae and zooxanthellae which inhabit hydras, infusoria, etc.; that there are organisms, (e. g., the lower Cyanophyceae, such as *Aphanocapsis* and *Microcystis*) which can be considered as free-living chromatophores; that certain Cyanophyceae actually live as symbionts in the cell plasma. This theory he thinks, is the only possible explanation of the polyphyletic origin of primeval plants, which were merely amoebae and flagellates into which Cyanophyceae migrated; that the green, red, and brown Cyanophyceae account for the algae of these colors; that the plant cell-wall is due to the formation, by the symbiotic chloroplasts, of carbohydrates easily polymerized into cellulose; which wall makes impossible the further taking of solid food and entails the quiescent nature and simple organization of plants, minus nerve, muscle, and psychic life. Here is another pyramid of theory resting on its apex.—C. R. B.

¹⁴GUTTENBERG, H. R., VON, Die Lichtsinnesorgane der Laubblätter von *Adoxa Moschatellina* und *Cynocrambe prostrata*. Ber. Deutsch. Bot. Gesells. **23**: 265-273. pls. 10, 11. 1905.

¹⁵MERESCHKOWSKY, C., Ueber Natur und Ursprung der Chromatophoren im Pflanzenreiche. Biol. Centralbl. **25**: 593-604. 1905.